

The generalized no-ghost theorem for N=2 SUSY critical strings¹

Jadwiga Bieńkowska

*Department of Physics and
Enrico Fermi Institute
the University of Chicago
Chicago, Il 60637*

Abstract

We prove the no-ghost theorem for the N=2 SUSY strings in (2,2) dimensional flat Minkowski space. We propose a generalization of this theorem for an arbitrary geometry of the N=2 SUSY string theory taking advantage of the N=4 SCA generators present in this model. Physical states are found to be the highest weight states of the N=4 SCA.

¹work supported in part by DOE grant DE-FG02-90ER40560.

The N=2 SUSY critical string theory with (2,2) real signature seems to be an interesting model to test duality properties of string theories [1]. It is particularly simple because the only degree of freedom in the model is believed to be a scalar field playing the role of the Kähler potential deformations. Even though there is no reason to suppose that there are other physical states present in this model there was up to date no explicit no-ghost theorem proven.

In this letter we present such a proof for the flat Minkowski space with (2,2) real signature. The proof is a direct generalization of the modern version of the no-ghost theorem by Thorn [6] which uses the BRST quantisation prescription [10]. We present the line of the proof for the N=2 critical string theory pointing out the differences between the N=0 [6] and N=2 case.

The N=2 string theory has one advantage over cases with lower N. It possesses an additional symmetry generated by the spectral flow [2]. In 4 real dimensions (critical N=2 strings) the spectral flow generators are dimension 1 operators and can be identified as currents of the local SU(2) (in Euclidean space) or SU(1,1) (in Minkowski type space) groups [3]. The enlarged set of generators of the N=2 theory includes all the N=4 SCA generators [4], with the gauge symmetry generators being only the ones belonging to the N=2 SCA subset. Using this larger set of the operators we can look at the states in this model as states from the Verma module of the N=4 SCA. We show that gauging away the N=2 SUSY is enough to prove that the only physical states are the highest weight states of the N=4 SCA. In the Euclidean model the only physical state is the identity, while in the Minkowski case there are enough states left in the physical subspace to make the model interesting.

The critical N=2 SUSY string lives in the two dimensional complex space with two complex bosons $a_n^{\mu(*)}$ and fermions $b_n^{\mu(*)}$ $\mu = 1, 2$ [5]. The free field representation of the N=2 SCA is:

$$\begin{aligned}
L_n &= -\sum_s a_{n-s}^* \cdot a_s + \sum_s \left(\frac{n}{2} - s\right) b_{n-s}^* \cdot b_s \\
G_n &= \sqrt{2} \sum_s b_{n-s}^* \cdot a_s \quad G_n^* = \sqrt{2} \sum_s b_{n-s} \cdot a_s^* \\
T_n &= -\frac{1}{2} \sum_s b_{n-s}^* \cdot b_s
\end{aligned} \tag{1}$$

where $[a_r^\mu, a_s^{nu}] = -r\eta^{\mu\nu}\delta_{r+s}$, $\{b_r^\mu, b_s^{nu}\} = -\eta^{\mu\nu}\delta_{r+s}$. Dots signify a scalar product with respect to the $\eta^{\mu\nu}$ Minkowski ($diag(1, -1)$) or Euclidean ($diag(1, 1)$) hermitian

metric. The generators obey the N=2 SCA comutation relations:

$$\begin{aligned}
[L_n, L_m] &= (n-m)L_{n+m} + \frac{1}{2}kn(n^2-1)\delta_{n+m} \\
\{G_r, G_s^*\} &= 2L_{r+s} + 2(r-s)T_{r+s} + \frac{1}{2}k(4r^2-1)\delta_{r+s} \\
[L_m, G_r] &= (\frac{m}{2}-r)G_{m+r} \quad [L_m, G_r^*] = (\frac{m}{2}-r)G_{m+r}^* \\
[T_m, G_r] &= \frac{1}{2}G_{m+r} \quad [T_m, G_r^*] = -\frac{1}{2}G_{m+r}^* \\
[L_m, T_n] &= -nT_{m+r} \quad [T_m, T_n] = \frac{1}{2}km\delta_{n+m}
\end{aligned} \tag{2}$$

and all other commutators are zero . The moding of $b_r^{\mu(*)}, G_r^{(*)}$ operators is governed by the spectral flow parameter η [2],[5] and we will restrict our attention to one sector, let us say the NS with $r \in \mathbf{Z} + \frac{1}{2}$. The physical states are defined by the conditions

$$L_n|s\rangle = G_r|s\rangle = G_r^*|s\rangle = T_n|s\rangle = 0 \text{ for } n, r > 0 \quad L_0|s\rangle = T_0|s\rangle = 0 \tag{3}$$

In the BRST formulation of the theory for each gauge constraint one introduces a pair of ghosts with a statistics opposite to that of the constraint [5],[10]. For N=2 there are four pairs of ghosts, associated with L_n, T_n, G_n, G_n^* gauge constraints respectively, satisfying the commutation relations $\{c_n, \bar{c}_m\} = \delta_{m+n}$, $\{t_n, \bar{t}_m\} = \delta_{m+n}$, $[\gamma_n, \bar{\gamma}_m] = \delta_{m+n}$, $[\gamma_n^*, \bar{\gamma}_m^*] = \delta_{m+n}$. The BRST operator Q expressed in terms of these operators is

$$\begin{aligned}
Q &= L_n c_{-n} + G_p \gamma_{-p} + G_p^* \gamma_{-p}^* + T_n t_{-n} - \frac{1}{2}(m-n)c_{-m}c_{-n}\bar{c}_{n+m} - 2\gamma_{-p}\gamma_{-q}^*\bar{c}_{p+q} \\
&- 2(p-q)\gamma_{-p}\gamma_{-q}^*\bar{t}_{p+q} + nc_{-m}t_{-n}\bar{t}_{m+n} + (\frac{m}{2}-p)c_{-m}(\gamma_{-p}\bar{\gamma}_{m+p} + \gamma_{-p}^*\bar{\gamma}_{m+p}^*) \\
&+ \frac{1}{2}t_{-m}(\gamma_{-p}\bar{\gamma}_{m+p} - \gamma_{-p}^*\bar{\gamma}_{m+p}^*)
\end{aligned} \tag{4}$$

$Q^2 = 0$ for two complex bosons and fermions [5].

We are set to begin the proof of the no ghost theorem for the Minkowski signature N=2 SUSY critical string theory. We follow closely the line of the proof proposed by Thorn [6]. In Euclidean space the $L_0|state\rangle = 0$ condition trivially forces the only physical state to be the identity. To fix the gauge conditions (3) we introduce the gauge fixing operators $A_n^+, B_n^+, A_n^{*-}, B_n^{*-}$ where the notation $(+, -)$ is unambiguous in the (1,1) signature and $A_n^{\pm(*)}(B_r^{\pm(*)})$ refer to the free boson (fermion) operators projected on the lightlike directions. Following the line of the proof [6] we want to construct the Hilbert space of this model using gauge generators, gauge fixing

conditions and states which are simultaneously annihilated by all of them. We consider the states

$$\begin{aligned} & L_{-1}^{\lambda_1} \dots L_{-l}^{\lambda_l} G_{-1}^{g_1} \dots G_{-r}^{g_r} G_{-1}^{*g_1^*} \dots G_{-r^*}^{*g_r^*} T_{-1}^{\tau_1} \dots T_{-n}^{\tau_n} \\ & A_{-1}^{+\alpha_1} \dots A_{-a}^{+\alpha_a} B_{-1}^{+\beta_1} \dots B_{-b}^{+\beta_b} A_{-1}^{*-\alpha_1^*} \dots A_{-a^*}^{*-\alpha_a^*} B_{-1}^{*-\beta_1^*} \dots B_{-b^*}^{*-\beta_b^*} |t\rangle \end{aligned} \quad (5)$$

where $L_n|t\rangle = G_r|t\rangle = G_r^*|t\rangle = T_n|t\rangle = A_n^{*+}|t\rangle = B_r^{*+}|t\rangle = A_n^-|t\rangle = B_r^-|t\rangle = 0$ for $n, r > 0$ and $L_0|t\rangle = h|t\rangle$. The "mixed" $(+, -)$ gauge fixing conditions are needed to assure the linear independence of the vectors (5), as we explain below.

The first step is to prove the linear independence of the vectors having the form (5) at each level N . First of all we notice that the number of generators in the basis (5) is the same as the number of generators in the Hilbert space spanned by the free boson and free fermion operators acting on the vacuum with a momentum p $|0, p\rangle$, so at each level the number of vectors from the set (5) is the same as the number of vectors from the free field representation of the Hilbert space which clearly forms a linearly independent set (a basis of the vector space at each mass level).

What remains to be proven then is that at each level N we can express any vector from the basis of $A_n^\pm, A_m^{*\pm}, B_r^\pm, B_s^{*\pm}$ by the vectors from (5). This is enough since at each level N we have a finite dimensional vector space. The proof follows by induction. For $N = \frac{1}{2}$ the check is trivial since the $G_{-\frac{1}{2}}, G_{-\frac{1}{2}}^*$ generators take the role of $B_{-\frac{1}{2}}^-, B_{-\frac{1}{2}}^{*+}$ operators. At level $N = 1$ we check by straightforward algebra that any vector from the free field basis can be expressed by vectors from the set (5). In particular at this level it becomes clear that we need the mixed gauge fixing conditions as above to ensure the linear independence of vectors (5). We require that both $A_0^\pm$ are different from zero. The case when either of them is equal zero is easy to prove since the physical state condition $L_0|s\rangle = 0$ automatically requires that there are no mass excitations different from 0. The $p = 0$ case has to be addressed separately and we discuss it at the end of the main line of the proof.

Having checked the validity of our statement at level 1, we assume that it is true for any vector at level $n < N$, i.e. states from free field basis at mass level n can be expressed in terms of vectors (5). Any state from the free field basis at level N which is a product of several $A_n^{\pm(*)}(B_r^{\pm(*)})$ operators can be expressed in the basis (5) by the induction assumption and using the commutation relations with the gauge generators. The $B_{-N+\frac{1}{2}}^{*+}|h\rangle, B_{-N+\frac{1}{2}}^-|h\rangle$ states are expressed easily by $G_{-N+\frac{1}{2}}|h\rangle = A_0^- B_{-N+\frac{1}{2}}^{*+}|h\rangle + \text{other states from the basis}$,

$G_{-N+\frac{1}{2}}^*|h\rangle = A_0^{*+}B_{-N+\frac{1}{2}}^-|h\rangle + \text{other states from the basis}$. The states $A_{-N}^{*+}|h\rangle, A_{-N}^-|h\rangle$ can also be expressed in this basis as follows. Consider the expansion of the state $G_{-p}G_{-N+p}^*|h\rangle$ ($\frac{1}{2} < p < N - \frac{1}{2}$ and $p \in \mathbf{Z} + \frac{1}{2}$) in the free fields basis. We see that it contains the states $(A_0^{*-}A_{-N}^+ + A_0^{*+}A_{-N}^-)|h\rangle$ plus other states including the product of the several operators from the free field basis. By induction we can express these states in the (5) basis with one exception. There is the possibility that doing so we will cancel the $A_0^{*+}A_{-N}^-|h\rangle$ term. The cancellation occurs from the state $2A_0^-A_0^{*+}B_{p-N}^-B_{-p}^{*+}|h\rangle$ if we try to express the B_{-p}^{*+} operator using the G_{-p} gauge generator and then commute the B_{-N+p}^- to the right. To solve this problem we realize that $T_{-N} = -\frac{1}{2}\sum_{p=\frac{1}{2}}^{N-\frac{1}{2}}(B_{-p}^{*+}B_{p-N}^- + B_{-p}^{*+}B_{p-N}^-)|h\rangle$ and so the state $\sum_{p=\frac{1}{2}}^{N-\frac{1}{2}}B_{-p}^{*+}B_{p-N}^-$ can be expressed in the basis as shown above. Then it follows from $(\sum_{p=\frac{1}{2}}^{N-\frac{1}{2}}G_{-p}G_{-N+p}^* + 4A_0^-A_0^{*+}T_{-N})|h\rangle = -2NA_0^{*+}A_{-N}^- + \text{other states}$ where now all *other states* are from the basis (5), that we can express A_{-N}^- in this basis. The remaining state $A_{-N}^{*+}|h\rangle$ can then be expressed by expanding $L_{-N}|h\rangle = A_0^-A_{-N}^{*+}|h\rangle + \text{other states}$ where all *other states* can be expressed in the basis (5). This completes the proof of the linear independence of the vectors from the set (5).

The proof of the no-ghost theorem is now relatively simple if we follow Thorn's approach [6]. We consider the BRST Hilbert space built as follows from ghosts and vectors (5). We introduce the N=2 SCA generators which include the ghosts operators [5]:

$$\begin{aligned} \{Q, \bar{c}_n\} &= L_n + L_n^{gh} = \mathcal{L}_n & \{Q, \bar{t}_n\} &= T_n + T_n^{gh} = \mathcal{T}_n \\ [Q, \bar{\gamma}_p] &= G_p + G_p^{gh} = \mathcal{G}_p & [Q, \bar{\gamma}_p^*] &= G_p^* + G_p^{*gh} = \mathcal{G}_p^* \end{aligned} \quad (6)$$

We define the ghost vacuum as $\bar{c}_0|0\rangle_{ghost} = \bar{t}_0|0\rangle_{ghost} = 0$. Then our BRST basis is formed by the states

$$\begin{aligned} &\mathcal{L}_{-1}^{\lambda_1} \dots \mathcal{L}_{-l}^{\lambda_l} \bar{c}_{-1}^{\bar{\kappa}_1} \dots \bar{c}_{-k}^{\bar{\kappa}_k} \mathcal{G}_{-1}^{g_1} \dots \mathcal{G}_{-r}^{g_r} \bar{\gamma}_{-1}^{\bar{\chi}_1} \dots \bar{\gamma}_{-s}^{\bar{\chi}_s} \\ &\mathcal{G}_{-1}^{*g_1^*} \dots \mathcal{G}_{-r^*}^{*g_{r^*}^*} \bar{\gamma}_{-1}^{*\bar{\chi}_1^*} \dots \bar{\gamma}_{-s^*}^{*\bar{\chi}_{s^*}^*} \mathcal{T}_{-1}^{\tau_1} \dots \mathcal{T}_{-n}^{\tau_n} \bar{t}_{-1}^{\bar{\nu}_1} \dots \bar{t}_{-m}^{\bar{\nu}_m} \\ &A_{-1}^{+\alpha_1} \dots A_{-a}^{+\alpha_a} c_{-1}^{\kappa_1} \dots c_{-k}^{\kappa_k} B_{-1}^{+\beta_1} \dots B_{-b}^{+\beta_b} \gamma_{-1}^{\chi_1} \dots \gamma_{-s}^{\chi_s} \\ &A_{-1}^{*-\alpha_1^*} \dots A_{-a^*}^{*-\alpha_{a^*}^*} t_{-1}^{\kappa_1^*} \dots t_{-k^*}^{\kappa_{k^*}^*} B_{-1}^{*-\beta_1^*} \dots B_{-b^*}^{*-\beta_{b^*}^*} \gamma_{-1}^{*\chi_1^*} \dots \gamma_{-s^*}^{*\chi_{s^*}^*} |t\rangle_{gh} \end{aligned} \quad (7)$$

where $|t\rangle_{gh}$ means now a tensor product of the ghost vacuum and previously defined $|t\rangle$ states from the physical sector. The above vectors form a linearly independent set since adding ghosts does not destroy the linear independence of the basis (5). A

more useful for the no-ghost theorem is the basis in which we symmetrise the bosonic operators and antisymmetrise the fermionic ones. Using the N=2 SCA commutation rules (2) and the ghosts commutation properties we can do it without spoiling the linear independence of the basis (7). We introduce shorthand notation to refer to this symmetrised basis

$$\begin{aligned} & \mathcal{L}_{\{-l\}}^\lambda \quad \bar{c}_{[-\bar{k}]}^{\bar{\kappa}} \mathcal{G}_{[-r]}^g \bar{\gamma}_{\{-s\}}^{\bar{\chi}} \mathcal{G}_{[-r^*]}^{*g^*} \bar{\gamma}_{\{-s^*\}}^{*\bar{\chi}^*} \mathcal{T}_{\{-n\}}^\tau \bar{t}_{[-m]}^{\bar{\nu}} \\ & A_{\{-a\}}^{+\alpha} c_{[-k]}^\kappa B_{[-b]}^{+\beta} \gamma_{\{-s\}}^\chi A_{\{-a^*\}}^{*-\alpha^*} t_{[-k^*]}^{\kappa^*} B_{[-b^*]}^{*-\beta^*} \gamma_{\{-s^*\}}^{*\chi^*} |t\rangle_{gh} \end{aligned} \quad (8)$$

where the $\{\}, []$ means symmetrization or antisymmetrization in the indices enclosed.

Following the proof by Thorn [6] we can easily verify that the specific symmetry combination of the first pair $\mathcal{L}_{-n}, \bar{c}_{-m}$ can be expressed by other states with a lower number of $\mathcal{L}$'s and a pure gauge. Working down the number of $\mathcal{L}$'s we can express any state with a symmetry associated to the Young tableau $\begin{array}{c} \lambda \\ \bar{\kappa} + 1 \end{array} \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array}$ by other states

and a pure gauge. Thus we are left with the states of the symmetry $\begin{array}{c} \lambda + 1 \\ \bar{\kappa} \end{array} \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array}$ only. Further requirement that these states be physical (annihilated by Q) drives us to the conclusion that the symmetry properties exclude any state with $(\lambda, \bar{\kappa}) \neq (0, 0)$ to be physical (for a more detailed discussion see [6]).

This procedure can be repeated smoothly for all other pairs of gauge generators and antighosts. We are left with the basis formed by the gauge fixing operators and ghosts $A_{\{-a\}}^{+\alpha} c_{[-k]}^\kappa B_{[-b]}^{+\beta} \gamma_{\{-s\}}^\chi A_{\{-a^*\}}^{*-\alpha^*} t_{[-k^*]}^{\kappa^*} B_{[-b^*]}^{*-\beta^*} \gamma_{\{-s^*\}}^{*\chi^*} |t\rangle_{gh}$. We can easily see from the commutation relations

$$\begin{aligned} [Q, A_{-m}^+] &= mc_{-n} A_{n-m}^+ + m\sqrt{2}\gamma_{-n}^* B_{n-m}^+ \\ [Q, A_{-m}^{*-}] &= mc_{-n} A_{n-m}^{*-} + m\sqrt{2}\gamma_{-n} B_{n-m}^{*-} \\ \{Q, B_{-m}^+\} &= (m - \frac{n}{2})c_{-n} B_{n-m}^+ + \frac{1}{2}t_{-n} B_{n-m}^+ + \sqrt{2}\gamma_{-n} A_{n-m}^+ \\ \{Q, B_{-m}^{*-}\} &= (m - \frac{n}{2})c_{-n} B_{n-m}^{*-} + \frac{1}{2}t_{-n} B_{n-m}^{*-} + \sqrt{2}\gamma_{-n} A_{n-m}^{*-} \end{aligned} \quad (9)$$

that the procedure of removing specific symmetry combinations of gauge fixing conditions and ghosts works when $A_0^\pm \neq 0$. We can remove, in the standard way [6], the first pair $A_{\{-a\}}^{+\alpha} c_{[-k]}^\kappa$. Then by the BRST charge Q acting on the vector formed from the basis $B_{[-b]}^{+\beta} \gamma_{\{-s\}}^\chi B_{[-b^*]}^{*-\beta^*} \gamma_{\{-s^*\}}^{*\chi^*} A_{\{-a^*\}}^{*-\alpha^*} t_{[-k^*]}^{\kappa^*} |t\rangle_{gh}$ we will reproduce, due to the commutation relations (9), the states containing operators A_{-n}^+, c_{-m} [6]. By the

previous step, since we consider the states annihilated by Q , states containing any of these operators have to be a pure gauge. The procedure of removing the states with $(\beta, \chi) \neq (0, 0)$ and $(\beta^*, \chi^*) \neq (0, 0)$ from this basis follows.

We are left with the states formed from the basis $A_{\{-a^*\}}^{*-a^*} t_{[-k^*]}^{\kappa^*} |t\rangle_{gh}$. We require that for a linear combination of the above states $|s\rangle$ $Q|s\rangle = 0$. Looking at the commutation relations (9) and $[Q, t_{-n}] = mc_{m-n}t_{-m} - 2(m-2n)\gamma_{m-n}\gamma_{-m}^*$ we see that it is enough to consider only the γ_{-p}^* ghost conservation number. The states with the non zero γ_{-p}^* ghost have to cancel among themselves and starting from the state with the highest t_{-n} excitation we see that the only physical state $Q|phys\rangle = 0$ is the one with no t ghost. Then it is easy to see that the states built only from A_{-n}^{*-} excitations are not physical. This completes the proof for $A^\pm \neq 0$.

For $A^+ = 0$ or $A^- = 0$ the physical state condition $L_0|phys\rangle = 0$ automatically excludes any excitation so we are left with the physical states subspace $|0, p\rangle$ where $p^\mu p_\mu^* = 0$.

The $p^\mu = 0$ case is special, as we know from the N=0,1 SUSY case [7]. We have to check directly which states with zero L_0, T_0 eigenvalue are physical. In this case $c_1|0\rangle_{gh} \neq 0$ ($L_{-1}|0\rangle = 0$), $\gamma_{\frac{1}{2}}|0\rangle_{gh} \neq 0$ ($G_{-\frac{1}{2}}|0\rangle = 0$), $\gamma_{\frac{1}{2}}^*|0\rangle_{gh} \neq 0$ ($G_{-\frac{1}{2}}^*|0\rangle = 0$) and there are several nontrivial possibilities for the states with non zero ghost numbers to be physical $L_0|s\rangle = 0$. We investigated all possible states built up from the above ghosts and we found that there are in this case additional non zero ghost number physical states

$$(a_{-1}^\mu c_1 + \sqrt{2}b_{-\frac{1}{2}}^\mu \gamma_{\frac{1}{2}}^*)|0, p^\mu = 0\rangle \quad (a_{-1}^{*\mu} c_1 + \sqrt{2}b_{-\frac{1}{2}}^{*\mu} \gamma_{\frac{1}{2}})|0, p^\mu = 0\rangle \quad (10)$$

This completes the proof for the no-ghost theorem for the N=2 SUSY critical strings in flat space.

We generalize the no-ghost theorem for the case of an arbitrary geometry of the N=2 SUSY string theory in (2,2) signature real dimension space by taking advantage of the larger set of generators existing in these theories due to the spectral flow [2], [3]. In the case of the (4,0) real signature space the group is enlarged by the spectral flow generators T^+, T^- which can be identified with the generators of the SU(2) group. The full set of generators is described by the N=4 SCA operators [3], [4]. One can check that in the free field representation (1) the N=2 SCA algebra is completed to the N=4 by adding the generators.

$$T_n^+ = \sum_s b_{n-s}^{*1} b_s^{*2} \quad T_n^- = - \sum_s b_{n-s}^1 b_s^2 \quad (11)$$

It is important to have a hermitian conjugate basis for the operators $(T_n^+)^\dagger = T_n^-$ since this unambiguously defines the commutation relations in the algebra. The additional generators of the N=4 SCA and their commutation rules (non-zero ones) are

$$\begin{aligned}
[T_n^+, G_r^*] &= \sqrt{2} \sum_t (b_{n+r-t}^{*2} a_t^{*1} - b_{n+r-t}^{*1} a_t^{*1}) = B_{n+r} \\
[T_n^-, G_r] &= -\sqrt{2} \sum_t (b_{n+r-t}^2 a_t^1 - b_{n+r-t}^1 a_t^1) = -B_{n+r}^* \\
[T_n^+, T_m^-] &= 2T_{n+m} + \frac{1}{2}kn\delta_{n+m} \quad [T_n, T_m^\pm] = \pm T_{m+n}^\pm \\
[T_m^+, B_r^*] &= -G_{r+m} \quad [T_m^-, B_r] = G_{r+m}^* \\
\{B_r, B_s^*\} &= 2L_{r+s} + 2(r-s)T_{r+s} + \frac{1}{2}k(4r^2 - 1)\delta_{r+s} \\
[T_m, B_r] &= \frac{1}{2}B_{r+m} \quad [T_m, B_r^*] = -\frac{1}{2}B_{r+m}^* \\
\{B_r, G_s\} &= 2(r-s)T_{s+r}^+ \quad \{B_r^*, G_s^*\} = -2(r-s)T_{r+s}^-
\end{aligned} \tag{12}$$

which is equivalent to the N=4 SCA with the SU(2) subgroup discussed in [4] with $G_r = \frac{1}{\sqrt{2}}(G_r^1 + \bar{G}_r^2)$, $G_r^* = \frac{1}{\sqrt{2}}(G_r^2 + \bar{G}_r^1)$, $B_r = \frac{1}{\sqrt{2}}(-G_r^1 + \bar{G}_r^2)$, $B_r^* = \frac{1}{\sqrt{2}}(G_r^2 - \bar{G}_r^1)$. The above defined operators are hermitian in the sense $(B_r)^\dagger = B_{-r}^*$, $(G_r)^\dagger = G_{-r}^*$. It is then straightforward to find the N=4 SCA in the case of Minkowski signature space. The algebra has, as could be expected, the SU(1,1) subgroup and the commutations relations for the generators $T_n^\pm$ and $B_r^{(*)}$ defined as in (11, 12):

$$\begin{aligned}
[T_n^+, T_m^-] &= -2T_{n+m} + \frac{1}{2}kn\delta_{n+m} \\
[T_m^+, B_r^*] &= G_{r+m} \quad [T_m^-, B_r] = -G_{r+m}^* \\
\{B_r, B_s^*\} &= -(2L_{r+s} + 2(r-s)T_{r+s}) - \frac{1}{2}k(4r^2 - 1)\delta_{r+s}
\end{aligned} \tag{13}$$

where we have listed only the commutators which are different from case in (12) .

The algebra we obtain in this case is considerably different from the Euclidean space algebra, since it contains as a subgroup the noncompact SU(1,1). The SU(1,1) group is the part of the rotation group unbroken by the background. Typical physical states of the theory belong to finite dimensional representations of SU(1,1). Fortunately the representations of this group were classified some time ago [8]. There are three basic types of the SU(1,1) representations classified by the eigenvalues of the quadratic Casimir operator $C = -g_{ab}T_0^a T_0^b$, where $g_{ab} = \text{diag}(-1, -1, 1)$

$C|0, j, m\rangle = -j(j+1)|0, j, m\rangle$, and the discrete values of the compact direction operator $T_0^3|0, j, m\rangle = T_0|0, j, m\rangle = m|0, j, m\rangle$. The ground states of the N=4 SCA theory (13) belong to the irreducible representations of the global SU(1,1) algebra and we introduced notation $|0, j, m\rangle$ where 0 refers to the state zero excitation of the N=4 SCA. There are following irreducible (non necessarily unitary) SU(1,1) representations: continuous ones

- 1) C_j^0 , $j = -\frac{1}{2} + i\kappa$ and $m = 0, \pm 1, \pm 2 \dots$
- 2) $C_j^{\frac{1}{2}}$, $j = -\frac{1}{2} + i\kappa$ and $m = \pm \frac{1}{2}, \pm \frac{3}{2} \dots$
- 3) E_j , $j \in \mathbf{R}$, $j \notin \mathbf{Z}$, $2j \notin \mathbf{Z}$ $m = 0, \pm 1, \pm 2, \dots$ or $m = \pm \frac{1}{2}, \pm \frac{3}{2}, \dots$

and there is no highest state annihilated either by T_0^+ or T_0^- ; infinite discrete representations:

- 4) D_j^+ , $j = -\frac{1}{2}, -1, -\frac{3}{2} \dots$ and $m = -j, -j+1, \dots$
- 5) D_j^- , $j = -\frac{1}{2}, -1, -\frac{3}{2} \dots$ and $m = j, j-1, \dots$

and finite discrete representations;

- 6) F_j , $j = \frac{1}{2}, 1, \dots$ and $m = -j, -j+1, \dots, j-1, j$

For the D_j^+ representation there exists a highest state $|s\rangle^+ = |0, j-j\rangle$ such that $T_0^-|s\rangle^+ = 0$ and for the D_j^- representation there is a state $|s\rangle^- = |0, j, j\rangle$ such that $T_0^+|s\rangle^- = 0$. These remarks are useful in the generalization of the no-ghost theorem we present below.

The space of all states in the theory we consider is a tensor product of the states of the Verma module of the N=4 SCA, obeying the commutation relations (2, 12, 13) (SU(1,1) case), and some other states which are the trivial representation of the N=4 SCA. The goal of our proof is to show that requirement that the states obey the N=2 SCA gauge conditions (3) is enough to gauge away all excitations from the N=4 SCA Verma module. This leaves the physical states space of the theory containing only the highest weight states of the N=4 SCA.

We consider a state from a Verma module built on a state belonging to one of the global representations of the SU(1,1) algebra. The states built from the ordered product of the N=4 SCA operators acting on the highest weight state:

$$\begin{aligned} & L_{-1}^{\lambda_1} \dots L_{-l}^{\lambda_l} G_{-1}^{g_1} \dots G_{-r}^{g_r} G_{-1}^{g_1^*} \dots G_{-r^*}^{g_r^*} T_{-1}^{\tau_1} \dots T_{-n}^{\tau_n} \\ & T_{-1}^{+\alpha_1} \dots T_{-a}^{+\alpha_a} B_{-1}^{\beta_1} \dots B_{-b}^{\beta_b} T_{-1}^{*\alpha_1^*} \dots T_{-a^*}^{*\alpha_a^*} B_{-1}^{*\beta_1^*} \dots B_{-b^*}^{*\beta_b^*} |0, j, m\rangle \end{aligned} \quad (14)$$

form a good basis for the Verma module, in the sense that there are no linear combinations of them equaling zero (of course there are linear combinations of states with

norm equal to zero). The ordering of the operators (last four $T_{-n}^+, B_{-p}, T_{-n}^-, B_{-p}^*$) is important for the no-ghost proof and should be set respectively to the representation of the $|0, j, m\rangle$ state. For the continuous representations $C_j^{\frac{1}{2}}, C_j^0, E_j$ it does not matter, for the D_j^- representations the positive charge generators T_m^+, B_n should come to the left as in (14) and for the D_j^+ representations the negative charge operators T_m^-, B_n^* come first from the left.

The no ghost theorem follows in much the same way as for flat space. However there are some differences which we point out below. We also restrict our discussion to the case when the ground state belongs to D_j^- or continuous representation, but the D_j^+ case is proven analogously.


To gauge away the N=2 SUSY we use the BRST prescription with the BRST charge defined by (4). The full BRST Hilbert space takes the familiar form

$$\begin{aligned} \mathcal{L}_{\{-l\}}^\lambda \quad \bar{c}_{[-\bar{k}]}^\kappa \quad \mathcal{G}_{[-r]}^g \quad \bar{\gamma}_{\{-\bar{s}\}}^{\bar{\chi}} \quad \mathcal{G}_{[-r^*]}^{g^*} \quad \bar{\gamma}_{\{-\bar{s}^*\}}^{*\bar{\chi}} \quad \mathcal{T}_{\{-n\}}^\tau \quad \bar{t}_{[-\bar{m}]}^\nu \\ T_{\{-a\}}^{+\alpha} \quad c_{[-k]}^\kappa \quad B_{[-b]}^\beta \quad \gamma_{\{-s\}}^\chi \quad T_{\{-a^*\}}^{-\alpha^*} \quad t_{[-k^*]}^{\kappa^*} \quad B_{[-b^*]}^{\beta^*} \quad \gamma_{\{-s^*\}}^{*\chi^*} \quad |0, j, m\rangle_{gh} \end{aligned} \quad (15)$$

Adding ghosts and the (anti) symmetrization procedure does not spoil the linear independence of the basis (14). But unlike in the previous case the subspace $\mathcal{H}_{TB}$ built from the vectors $T_{\{-a\}}^{+\alpha} c_{[-k]}^\kappa B_{[-b]}^\beta \gamma_{\{-s\}}^\chi T_{\{-a^*\}}^{-\alpha^*} t_{[-k^*]}^{\kappa^*} B_{[-b^*]}^{\beta^*} \gamma_{\{-s^*\}}^{*\chi^*} |0, j, m\rangle_{gh}$ is not Q invariant. There exist vectors $|s\rangle \in \mathcal{H}_{TB}$ such that $Q|s\rangle \notin \mathcal{H}_{TB}$. This is the consequence of the commutation relations of these generators with the BRST charge

$$\begin{aligned} [Q, T_{-m}^+] &= mc_{-n} T_{n-m}^+ + T_{n-m}^+ t_{-n} - \gamma_{-n}^* B_{n-m} \\ [Q, T_{-m}^-] &= mc_{-n} T_{n-m}^- - T_{n-m}^- t_{-n} + \gamma_{-n} B_{n-m}^* \\ \{Q, B_{-m}\} &= -(m + \frac{n}{2}) B_{n-m} c_{-n} - \frac{1}{2} B_{n-m} t_{-n} - 2(n+m) \gamma_{-n} T_{n-m}^+ \\ \{Q, B_{-m}^*\} &= -(m + \frac{n}{2}) B_{n-m}^* c_{-n} + \frac{1}{2} B_{n-m}^* t_{-n} + (n+m) \gamma_{-n} T_{n-m}^- \end{aligned} \quad (16)$$

and the N=4 SCA (12), (13).

As in the previous case, we would like to express the states with a symmetry λ  in the $\mathcal{L}_{\{l\}}^\lambda$ and $\bar{c}_{[\bar{k}]}^\kappa$ operators successively by the states with the lower number of $\mathcal{L}_{-n}$'s and pure gauge terms and eventually eliminate them from the physical states space. In the case of the basis (15) there will be some states with the same number λ of $\mathcal{L}_{-n}$ representing the above symmetry state in our induction procedure [6], which did not happen previously. The unwanted $\mathcal{L}_{-n}$ generators will come from

the commutation relations (16) and a subsequent commutation of the B, B^* generators. We are getting one $\mathcal{L}_{-n}$ generator at the expense of exchanging two B, B^* pair for the ghost. For each finite N level space we can subsequently work down the number of $\mathcal{L}_{-n}$ -producing generators eliminating them after a finite number of steps, and so our procedure of eliminating the symmetry combination

$$\begin{array}{c} \lambda \\ \bar{\kappa} + 1 \\ 1 \end{array} \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} \text{ survives.}$$

Then we consider how the states with the remaining symmetry

$$\begin{array}{c} \lambda + 1 \\ \bar{\kappa} \end{array} \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array}$$

can be made physical and we come to the conclusion that the only possibility is that $(\lambda, \bar{\kappa}) = (0, 0)$. Working further with the basis of the states without the $\mathcal{L}$'s or $\bar{c}$'s we would again encounter the $\mathcal{L}$ state reproduced when we try to express the combination

$$^g \begin{array}{c} \bar{\kappa} + 1 \\ \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} \end{array}$$

by other states and pure gauge terms. But by the first step, and physical state condition, any such state has to be pure gauge and the procedure follows all the way thus taking care of all the states containing the gauge generators and antighosts.

The next difference showing up in this case is that the role of the A_0^+, A_0^{*-} operators (which are **C**-numbers in flat Minkowski space) is played by the $T_0^\pm$ as we see from (16). Since the ground states in this theory belong to the SU(1,1) infinite representations, this step of the proof also survives. To check it we need the basis ordered as was discussed before.

Looking at the commutator

$$\begin{aligned} [Q, T_{-\{a\}}^{+(\alpha^++1)} c_{-[k]}^{(\kappa-1)}] &= \sum_i T_{\{-1\}}^{+\alpha_1^+} \dots \hat{T}_{-i}^{+\alpha_i^+} \dots T_{-a\}^{+\alpha_a^+} i T_0^+ c_{-i} c_{-[k]}^{\kappa-1} + \\ &+ \text{ other states} \end{aligned} \quad (17)$$

we see that the state with the symmetry

$$\begin{array}{c} \alpha + 1 \\ \kappa \end{array} \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array}$$

can be expressed by other states and a pure gauge if there always exists a state at zero level $|0, j, m-1\rangle$ such that $|0, j, m\rangle = T_0^+ |0, j, m-1\rangle$. This statement is always true in the continuous and D_j^- representation since we ordered our basis to assure that the states containing $T_n^+, c_n, B_m, \gamma_m$ operators can be always removed. We can also remove in this way the states built from the $T_n^-, t_n, B_m^*, \gamma_m^*$ operators as long as the state $|0, j, m\rangle$ is different from the highest weight state of the D_j^- global representation $|s\rangle^-$. For this state there does not exist $|\tilde{s}\rangle$ such that $T_0^- |\tilde{s}\rangle = |s\rangle^-$. We have to consider states built on the highest state $|s\rangle^-$ separately. Any other state from the basis (16), which

has $|0, j, m\rangle$ some other state from the D_j^- or a continuous representation is either nonphysical or a pure gauge.

We have to check directly which linear combinations of $T_{\{-a^*\}}^{-\alpha^*} t_{[-k^*]}^{\kappa^*} B_{[-b^*]}^{*\beta^*} \gamma_{\{-s^*\}}^{*\chi^*} |s\rangle^-$ states can be physical e.i. annihilated by Q . When Q acts on the linear product of the above states we can realise that the terms $(\dots)T_0^- c_{-n}|s\rangle^- = (\dots)c_{-n}|0, j, j-1\rangle$ have to cancel by themselves, by the ghost conservation number and orthogonality of the states in the $SU(1,1)$ representation D_j^- . This excludes the possibility of the T_{-n}^- excitations in the physical state. Let us look next at the commutators of Q with the remaining generators. From the rules (16), $[Q, t_{-n}] = mc_{m-n}t_{-m} - 2(m-2n)\gamma_{m-n}\gamma_{-m}^*$ and $[Q, \gamma_{-n}^*] = (\frac{3}{2}m-n)c_{-m}\gamma_{m-n}^* - \frac{1}{2}t_{-m}\gamma_{m-n}^*$ we see that the only terms containing the γ_{-p} ghost come from the commutators with the t_{-n} ghost and therefore have to cancel by themselves. Starting from the state containing t_{-n} with the highest n we can easily see that the only possibility is that there are no t_{-n} ghost excitations at all. We are finally left with the linear combination of states $B_{[-b^*]}^{*\beta^*} \gamma_{\{-s^*\}}^{*\chi^*} |s\rangle^-$, and by looking at the $[Q, B_{-p}^*]$ commutators, we see the states of the form $T_0^- \gamma_{-p}^* |s\rangle^-$ have to cancel each other. Starting from the state with highest excitation number of B_{-n}^* and working down the B_{-p}^* 's number we see that there is no possibility to cancel the last one ($[Q, \gamma_p^*]$ does not contain the T_0^-) and so the only solution to the physical state condition is the one where $\beta^* = 0$. It is straightforward to see that the remaining states built out of the γ_{-p}^* ghosts cannot be physical for nonzero ghost excitation. For the $|0, j, m\rangle$ belonging to the finite dimension representations F_j we can gauge away any state from the basis (15) with $m > -j$ as above. We have to consider only the linear combination of states $T_{\{-a^*\}}^{-\alpha^*} c_{[-k]}^{\kappa} B_{[-b]}^{\beta} \gamma_{\{-s\}}^{\chi} T_{\{-a^*\}}^{-\alpha^*} t_{[-k^*]}^{\kappa^*} B_{[-b^*]}^{*\beta^*} \gamma_{\{-s^*\}}^{*\chi^*} |0, j, -j\rangle$. We require that Q acting on such states is zero. From the commutators (16) we see that states $[\dots]T_0^+ c_{-n}|0, j, -j\rangle$ have to cancel among themselves and so $\alpha = 0$. We can further eliminate any B_{-p} excitations realising that $[\dots]T_0^+ c_{-n}|0, j, -j\rangle$ states have to cancel each other. The remaining states $T_{\{-a^*\}}^{-\alpha^*} t_{[-k^*]}^{\kappa^*} B_{[-b^*]}^{*\beta^*} \gamma_{\{-s^*\}}^{*\chi^*} |0, j, -j\rangle$ are gauged away as in equation (17) since we can always express $|0, j, -j\rangle = T_0^- |0, j, -j+1\rangle$. This completes the proof that the states from the Verma module of the N=4 SCA are either nonphysical or pure gauge in the N=2 SUSY theory. The only possible physical states are the highest weight states of the N=4 SCA which fall into the representations of the global $SU(1,1)$ group.

The physical state constraints (3) could tell us more about which of these states are physical. We can think of the Hilbert space of this theory as a product of the

SU(1,1) Kac-Moody algebra representation and the space which is trivial with respect to it. The $L_0|state\rangle = 0$ condition does not provide additional information. From the Sugawara construction for the SU(1,1) group [9] and the value of the central charge $c=6$ [3] we see that $L_0^{SU(1,1)}|0,j,m\rangle = -j(j+1)|0,j,m\rangle$ and easily we get $-j(j+1) < 0$. Since the SU(1,1) trivial part of the space remains undetermined we can not say which combination of states from the global SU(1,1) representations are allowed. The charge of the state is that of the SU(1,1) representation and the requirement $T_0|state\rangle = 0$ means that we should consider only the SU(1,1) states with zero charge. This excludes the discrete unitary representations $D_j^\pm$, $C_j^{\frac{1}{2}}$, and noninteger m E_j representations since we have $m \neq 0$ in these cases.

In the proof presented the Verma modules built from the trivial ($j = 0$) representation of the global SU(1,1) were not considered. In such a case the presented above line of the proof breaks down since both operators $T_0^\pm$ annihilate the ground state of the theory. This situation is similar to the zero momentum case in Minkowski space but here we are unable to determine, from general arguments, what is the L_0 eigenvalue of the $|0, j = 0, 0\rangle$ state since we can not say much about its SU(1,1) trivial part.

I would like to thank Emil Martinec for bringing this problem to my attention and for many enlightening discussions. I thank also Peter Freund and Tohru Eguchi for helpful remarks. This work is submitted in partial fulfillment of the requirements for a Ph.D. degree in physics at the University of Chicago.

References

- [1] H.Ooguri and C.Vafa, *Geometry of N=2 strings* EFI91/05, HUTP-91/A003. Mod. Phys. Lett. A5 (1990),1389.
- [2] A.Schiwimmer and N.Seiberg, Phys. Lett. B184 (1987), 191.
- [3] T.Eguchi and A.Taormina, Nucl. Phys B315 (1989), 193.
- [4] T.Eguchi and A.Taormina, Phys. Lett. B196 (1987), 75. Phys. Lett. B200 (1988), 315. Phys. Lett. B210 (1988), 125.
- [5] S.D.Mathur and T.Mukhi, Phys. Rev D36(2) (1987), 465.
- [6] C.Thorn, Nucl. Phys B286 (1987), 61.

- [7] I.B.Frenkel,H.Garland and G.J.Zukermann, Proc.Natl.Acad.Sci. USA (1986), pp 8440-8446.
- [8] V.Bargman, Ann. of Math. 48(3) (1947), 568.
- [9] L.J.Dixon, M.Peskin and J.Lykken, Nucl. Phys B325 (1989), 329. S.Hwang, Nucl. Phys B354 (1991), 100.
- [10] I.A.Batalin and G.A.Vilkoviski, Phys. Lett. B69 (1977), 309.